

Cambridge IGCSE[™]

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

1 6 8 6 1 0 2 9 8 4

CAMBRIDGE INTERNATIONAL MATHEMATICS

0607/41

Paper 4 (Extended)

May/June 2024

2 hours 15 minutes

You must answer on the question paper.

You will need: Geometrical instruments

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You should use a graphic display calculator where appropriate.
- You may use tracing paper.
- You must show all necessary working clearly and you will be given marks for correct methods, including sketches, even if your answer is incorrect.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- For π , use your calculator value.

INFORMATION

- The total mark for this paper is 120.
- The number of marks for each question or part question is shown in brackets [].

This document has 20 pages. Any blank pages are indicated.

Formula List

For the equation

$$ax^2 + bx + c = 0$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Curved surface area, A, of cylinder of radius r, height h.

$$A = 2\pi rh$$

Curved surface area, A, of cone of radius r, sloping edge l.

$$A = \pi r l$$

Curved surface area, A, of sphere of radius r.

$$A = 4\pi r^2$$

Volume, V, of pyramid, base area A, height h.

$$V = \frac{1}{3}Ah$$

Volume, V, of cylinder of radius r, height h.

$$V = \pi r^2 h$$

Volume, V, of cone of radius r, height h.

$$V = \frac{1}{3}\pi r^2 h$$

Volume, V, of sphere of radius r.

$$V = \frac{4}{3}\pi r^3$$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$Area = \frac{1}{2}bc\sin A$$

Answer all the questions.

1	(a)	Solve the equations.			
		(i) $3x-2=-14$			
				<i>x</i> =	[2]
		(ii) $7x + 11 = 26 - 3x$			
				<i>x</i> =	[2]
	(b)	Solve the simultaneous equation You must show all your working	S.		
			5x + 3y = -15 $3x + 5y = -17$		
			3x + 3y = -17		
				<i>x</i> =	
				<i>y</i> =	[4]
	(c)	Solve the inequality.			
		2x+1 > 9			
					[4]

2 (a) The heights, x cm, of 100 plants are shown in the table.

Height (x cm)	$0 < x \le 20$	$20 < x \leqslant 35$	$35 < x \leqslant 40$	$40 < x \le 60$	$60 < x \le 80$
Frequency	7	13	20	32	28

(i) Calculate an estimate of the mean height of the plants.


......cm [2]

(ii) (a) Complete the cumulative frequency table for the plants.

Height (x cm)	<i>x</i> ≤ 20	<i>x</i> ≤ 35	<i>x</i> ≤ 40	<i>x</i> ≤ 60	<i>x</i> ≤ 80
Cumulative frequency	7				100

[1]

(b) On the grid, draw the cumulative frequency curve.



[3]

-	· - /	T T	cumulative fi		4 -	C 1		C 41		
	α	I ICA VALIT	cumiliative ti	eallenev	CHIEVE IO	Tina an	ectimate	TOT THE	interdilarille	range
•	v	USC YOUI	Cumulan vC m	cquency	cui ve to	minu an	CStilliate	ioi uic	IIIICI quai tiic	range.


..... cm [2]

(b) The heights, $h \, \text{cm}$, of 50 different plants are shown in the table, where k is an integer.

Height (h cm)	Frequency
0 < h ≤ 20	25
$20 < h \leqslant k$	15
$k < h \leqslant 80$	10

An estimate	of the mean	height of the	ese plants is 27 c	m

Find the value of k.




(a)	Translate triangle A with vector $\begin{pmatrix} 2 \\ -6 \end{pmatrix}$. Label the image B.	[2]
(b)	Describe fully the single transformation that maps triangle B onto triangle A .	
		[2]
(c)	Rotate triangle A through 90° clockwise about $(0, 0)$. Label the image C .	[2]
(d)	Reflect triangle A in the line $y = x$. Label the image D.	[2]
(e)	Describe fully the single transformation that maps triangle C onto triangle B .	
		[3]

(a)	The price of a coat is \$84. The price is reduced by 12%.	
	Find the new price of the coat.	
		\$[2]
(b)	The price of a table is reduced by 25%. The price is now \$960.	
	Find the original price of the table.	
		\$[2]
(c)	Samir invests \$600 in a bank that pays compound interest	
	(i) Find the value of Samir's investment after 4 complet	te years.
		\$[2]
	(ii) Find the number of complete years for the value of more than \$1000.	of Samir's investment to be first worth
		[4]

(a)	Am	Amir and Bob work together and share their earnings in the ratio 3:5.						
	(i)	Find the amount Bob receives when their earnings are \$120.						
		\$	[2]					
	(ii)	They decide to change the ratio for all further earnings. Amir's share of the earnings is increased by 20% of his original share. Bob's share of the earnings is decreased by 20% of his original share.						

Show that the ratio of their earnings is now 9:10.

[3]



A, B, C and D lie on a circle, centre O. AP is a tangent to the circle at A. OP is perpendicular to AC and AD is parallel to BC. Angle $ABC = 61^{\circ}$ and angle $PAD = 22^{\circ}$.

(a)	Write down the mathematical	name of the cyclic	quadrilateral ABCD.
-----	-----------------------------	--------------------	---------------------

(b)	Complete the statement.	
	Angle $OAP = 90^{\circ}$ because	
		[1]

(c) Find

Angle
$$ADC = \dots$$
 [1]

.....[1]

(ii) angle ACD

(i) angle ADC


Angle
$$ACD = \dots$$
 [1]

(iii) angle ACB

Angle
$$ACB = \dots$$
 [2]

(iv) angle OCA.

Angle
$$OCA = \dots$$
 [2]



$$f(x) = 2 - \left| 1 - 0.5x^2 \right|$$

- (a) On the diagram, sketch the graph of y = f(x), for values of x between -3 and 3. [3]
- (b) The graph cuts the x-axis at points A and B.Work out the length AB.

$$AB = \dots$$
 [2]

(c) Solve f(x) = 0.5.

-[2]
- (d) Write down the coordinates of the minimum point of the graph.
 - (.....) [1]


(e) The equation f(x) = k has two solutions.

Find the range of values of k.

.....[2]


(a)	Spii	nner A and spinner B are each fair 5-sided spinners. nner A is numbered 1, 2, 2, 3, 4. nner B is numbered 1, 2, 3, 4, 4.		
	The	two spinners are each spun once and the number on each	ch spinner is recorded.	
	Fine	d the probability that		
	(i)	the number on spinner A is 6		
				[1]
	(ii)	the number on spinner B is not 4		
				[1]
	(iii)	the number on spinner A is the same as the number on	spinner B	
				[2]
	(iv)	the sum of the two numbers is 6.		[3]
	(11)	the sum of the two numbers is o.		
				[3]

(b) (i) On the Venn diagram, shade $A \cup B$.




[1]

(ii) Describe the shaded region using set notation.




.....[1]

(iii) The Venn diagram below shows the number of elements in each subset.



Find $n((A \cap B) \cap C')$.

.....[1]



The diagram shows a shape ABDC formed from triangle ABC and a sector of a circle BCD, centre B.


(a) Show that $BC = 9.0 \,\mathrm{m}$, correct to 1 decimal place.

[3]

(b) Use the sine rule to find angle BCA.

Angle
$$BCA = \dots$$
 [3]

(c)	Find the area of triangle <i>ABC</i> .	
(d)	Find the area of the shaded region.	m ² [2]
(e)	Find the perimeter of the shape <i>ABDC</i> .	m ² [3]
		m [2]



The diagram shows a solid cone with base radius 40 cm and slant height 60 cm.

(a) Find the volume of the cone.

cm ² [3		$m^3 [3]$
--------------------	--	-----------

(b) Show that the total surface area of the cone is $4000\pi\text{cm}^2$.


[2]

(c) A mathematically similar cone has a surface area of $1000\pi\text{cm}^2$.

Show that the radius of this cone is 20 cm.

[2]

(d)



A cone with radius 20 cm is removed from the top of the cone with radius 40 cm to leave a solid. Calculate the surface area of the remaining solid.

																						_					
								 					 						C	n	n			ſ	3	1	

10	y va $y =$	aries 18 v	inversely as the square root of $(x+1)$. when $x = 3$.		
	(a)	(i)	Find the value of y when $x = 8$.		
		(ii)	Find the value of x when $y = 1.5$.	<i>y</i> =	[3]
	(b)	w =	aries directly as the square root of $(x+1)$. = 18 when $x = 3$. If the value of \sqrt{wy} .	<i>x</i> =	[2]
				$\sqrt{wy} = \dots$	[3]

 $x \neq 1.5$

				1)
11		f(x) = 3x - 1	g(x) = 5 - 2x	$h(x) = \frac{1}{2x - 3},$
	(a)	Find f(4).		
	(b)	Solve $f(x) = -7$.		

$$g^{-1}(x) = \dots$$
 [2]

.....[1]

.....[2]

(d) Solve g(x) = 7h(f(x)). You must show all your working.

(c) Find $g^{-1}(x)$.

$$x =$$
 [6]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.